

Atlas of Genetics and Cytogenetics in Oncology and Haematology



OPEN ACCESS JOURNAL

INIST-CNRS

Leukaemia Section

Short Communication

t(9;15)(p13;q24) PAX5/GOLGA6A

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France (JLH)

Published in Atlas Database: June 2014

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/t0915p13q24ID1596.html>

DOI: 10.4267/2042/56302

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2015 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

Review on t(9;15)(p13;q24) PAX5/GOLGA6A, with data on clinics, and the genes implicated.

Clinics and pathology

Disease

B-cell acute lymphoblastic leukemia (B-ALL)

Note

Was not taken into account in this review a case of diffuse large B-cell lymphoma in a 51 year-old male patient (Morrison et al., 1994).

Epidemiology

One case to date, a 21-year-old male patient with a CD10+ (B-II, common) ALL (Coyaud et al., 2010).

Prognosis

No data.

Cytogenetics

Cytogenetics morphological

There were additional numerical anomalies (+5, +21).

Genes involved and proteins

PAX5

Location

9p13.2

Protein

391 amino acids; from N-term to C-term, PAX5 contains: a paired domain (aa: 16-142); an octapeptide (aa: 179-186); a partial homeodomain (aa: 228-254); a transactivation domain (aa: 304-359); and an inhibitory domain (aa: 359-391). Lineage-specific transcription factor; recognizes the consensus recognition sequence GNCCANTGAAGCGTGAC, where N is any nucleotide. Involved in B-cell differentiation. Entry of common lymphoid progenitors into the B cell lineage depends on E2A, EBF1, and PAX5; activates B-cell specific genes and repress genes involved in other lineage commitments.

Activates the surface cell receptor CD19 and repress FLT3.

Pax5 physically interacts with the RAG1/RAG2 complex, and removes the inhibitory signal of the lysine-9-methylated histone H3, and induces V-to-DJ rearrangements.

Genes repressed by PAX5 expression in early B cells are restored in their function in mature B cells and plasma cells, and PAX5 repressed (Fuxa et al., 2004; Johnson et al., 2004; Zhang et al., 2006; Cobaleda et al., 2007; Medvedovic et al., 2011).


GOLGA6A

Location

15q24.1

Protein

693 amino acids (aa); contains a coiled coil domain (aa 14-611), with 11 leucine zipper motifs (aa 360-430).



t(9;15)(p13;q24) PAX5/GOLGA6A (887 aa)

Jean Loup Huret 2014

© AtlasGeneticsOncology

PAX5/GOLGA6A fusion protein.

Member of the golgin family of proteins, localizes to the Golgi apparatus. Its function is poorly known (Gilles et al., 2000).

Result of the chromosomal anomaly

Hybrid gene

Description

Fusion of PAX5 exon 6 to GOLGA6A exon 3.

Fusion protein

Description

887 amino acids. The predicted fusion protein contains the DNA binding paired domain, the octapeptide, and the homeodomain of PAX5 and part of the coiled coil domain of GOLGA6A.

References

Morrison VA, Frizzera G, Arthur DC, Ogle KM, Hurd DD, Bloomfield CD, Peterson BA. Prognostic factors for therapeutic outcome of diffuse small non-cleaved cell lymphoma in adults. *Am J Hematol.* 1994 Aug;46(4):295-303

Gilles F, Goy A, Remache Y, Manova K, Zelenetz AD. Cloning and characterization of a Golgin-related gene from the large-scale polymorphism linked to the PML gene. *Genomics.* 2000 Dec 15;70(3):364-74

Fuxa M, Skok J, Souabni A, Salvagiotto G, Roldan E,

Busslinger M. Pax5 induces V-to-DJ rearrangements and locus contraction of the immunoglobulin heavy-chain gene. *Genes Dev.* 2004 Feb 15;18(4):411-22

Johnson K, Pflugh DL, Yu D, Hesslein DG, Lin KI, Bothwell AL, Thomas-Tikhonenko A, Schatz DG, Calame K. B cell-specific loss of histone 3 lysine 9 methylation in the V(H) locus depends on Pax5. *Nat Immunol.* 2004 Aug;5(8):853-61

Zhang Z, Espinoza CR, Yu Z, Stephan R, He T, Williams GS, Burrows PD, Hagman J, Feeney AJ, Cooper MD. Transcription factor Pax5 (BSAP) transactivates the RAG-mediated V(H)-to-DJ(H) rearrangement of immunoglobulin genes. *Nat Immunol.* 2006 Jun;7(6):616-24

Cobaleda C, Schebesta A, Delogu A, Busslinger M. Pax5: the guardian of B cell identity and function. *Nat Immunol.* 2007 May;8(5):463-70

Coyaud E, Struski S, Prade N, Familiades J, Eichner R, Quelen C, Bousquet M, Mugneret F, Talmant P, Pages MP, Lefebvre C, Penther D, Lippert E, Nadal N, Taviaux S, Poppe B, Luquet I, Baranger L, Eclache V, Radford I, Barin C, Mozziconacci MJ, Lafage-Pochitaloff M, Antoine-Poirel H, Charrin C, Perot C, Terre C, Brousset P, Dastugue N, Broccardo C. Wide diversity of PAX5 alterations in B-ALL: a Groupe Francophone de Cytopathologie Hematologique study. *Blood.* 2010 Apr 15;115(15):3089-97

Medvedovic J, Ebert A, Tagoh H, Busslinger M. Pax5: a master regulator of B cell development and leukemogenesis. *Adv Immunol.* 2011;111:179-206

This article should be referenced as such:

Huret JL. t(9;15)(p13;q24) PAX5/GOLGA6A. *Atlas Genet Cytogenet Oncol Haematol.* 2015; 19(2):148-149.